

### CLAIM AMENDMENTS

This listing of claims will replace all prior versions, and listings, of claims in the application:

Claims 1-75 (Canceled)

76. (Previously presented) A composition comprising a solid support coupled to a chemical group comprising a cleavable functional group, a tag and a reactive group covalently linked to a sample molecule, wherein said cleavable functional group is covalently coupled to said solid support, said tag is covalently coupled to said cleavable functional group, and said reactive group is covalently coupled to said tag and wherein said cleavable functional group, said tag and said reactive group are positioned relative to each other to allow transfer of said tag to said sample molecule and release of said sample molecule from said solid support upon cleavage of said cleavable functional group.

77. (Original) The composition of claim 76, wherein said sample molecule is selected from the group consisting of a polypeptide, a nucleic acid, a lipid, a second messenger, and a metabolite.

78. (Original) The composition of claim 77, wherein said sample molecule is a polypeptide.

79. (Original) The composition of claim 78, wherein said polypeptide has a modification selected from the group consisting of phosphorylation, glycosylation, ubiquitination, acetylation, palmitoylation, prenylation, sulfation, hydroxylation, and myristylation.

80. (Original) The composition of claim 79, wherein said polypeptide is a phosphopolypeptide.

81. (Original) The composition of claim 76, wherein the solid support is a glass bead.

82. (Original) The composition of claim 76, wherein said cleavable functional group is a chemical linker cleavable by light, an acid, a base or an enzyme.

83. (Original) The composition of claim 76, wherein said tag is a mass spectrometry tag.

84 (Original) The composition of claim 76, wherein said tag is selected from the group consisting of a stable isotope tag, an isotope distribution tag, and a charged amino acid.

85 (Original) The composition of claim 84, wherein said tag is a stable isotope coded amino acid.

86. (Original) The composition of claim 85, wherein said tag is a deuterated or non-deuterated amino acid.

87. (Withdrawn) The composition of claim 76, wherein said tag is a gas-phase basic group or a hydrophobic group.

88. (Withdrawn) The composition of claim 87, wherein the gas-phase basic group is pyridyl.

89. (Withdrawn) The method of claim 76, wherein said tag is selected from a fluorophore, chromophore, and spin label.

90. (Original) The composition of claim 76, wherein said tag comprises an element having a characteristic isotope distribution.

91. (Withdrawn) The composition of claim 90, wherein said element is chlorine or bromine.

92. (Original) The composition of claim 76, wherein said covalently linked reactive group is derived from a succinimide ester group or an iodoacetyl group.

93. (Original) The composition of claim 78, wherein a primary amine group of said polypeptide is modified by treatment with N-succinimidyl S-acetylthioacetate, hydroxylamine, and tris (2-carboxyethyl) phosphine.

94. (Previously presented) A composition comprising a solid support covalently coupled to a chemical group comprising a cleavable functional group, a mass spectrometry tag and a reactive group for covalently attaching a sample molecule, wherein said cleavable functional group is covalently coupled to said solid support, said tag is covalently coupled to said cleavable functional group, and said reactive group is covalently coupled to said tag and wherein said cleavable functional group, said tag and said reactive group are positioned relative to each other to allow transfer of said tag to a sample molecule attached to said reactive group upon cleavage of said cleavable functional group and release of said sample molecule from said solid support.

95. (Original) The composition of claim 94, wherein the solid support is a glass bead.

96. (Original) The composition of claim 94, wherein said cleavable functional group is a chemical linker cleavable by light, an acid, a base or an enzyme.

97. (Original) The composition of claim 94, wherein said tag is selected from the group consisting of a stable isotope tag, an isotope distribution tag, and a charged amino acid.

98. (Original) The composition of claim 97, wherein said tag is a stable isotope coded amino acid.

99. (Original) The composition of claim 98, wherein said tag is a deuterated or non-deuterated amino acid.

100. (Withdrawn) The composition of claim 94, wherein said tag is a gas-phase basic group or a hydrophobic group.

101. (Withdrawn) The composition of claim 100, wherein said gas-phase basic group is pyridyl.

102. (Withdrawn) The method of claim 94, wherein said tag is selected from a fluorophore, chromophore, and spin label.

103. (Original) The composition of claim 94, wherein said tag comprises an element having a characteristic isotope distribution.

104. (Withdrawn) The composition of claim 103, wherein said element is chlorine or bromine.

105. (Original) The composition of claim 94, wherein said reactive group of said chemical group is selected from the group consisting of a succinimide ester group and an iodoacetyl group.

106. (Previously presented) The composition of claim 76, wherein said cleavable group is a photocleavable group, said functional group is a mass tag, and said reactive group reacts with a sulfhydryl group.

107. (Previously presented) The composition of claim 106, wherein said sample molecule is selected from the group consisting of a polypeptide, a nucleic acid, a lipid, a second messenger, and a metabolite.

108. (Previously presented) The composition of claim 107, wherein said sample molecule is a polypeptide.

109. (Previously presented) The composition of claim 108, wherein said polypeptide has a modification selected from the group consisting of phosphorylation, glycosylation, ubiquitination, acetylation, palmitoylation, prenylation, sulfation, hydroxylation, and myristylation.

110. (Previously presented) The composition of claim 109, wherein said polypeptide is a phosphopolypeptide.

111. (Previously presented) The composition of claim 106, wherein the solid support is a glass bead.

112. (Previously presented) The composition of claim 106, wherein said mass tag is an amino acid.

113. (Previously presented) The composition of claim 112, wherein said mass tag is leucine.

114. (Previously presented) The composition of claim 106, wherein said mass tag is selected from the group consisting of a stable isotope tag, an isotope distribution tag, and a charged amino acid.

115. (Previously presented) The composition of claim 114, wherein said mass tag is a stable isotope coded amino acid.

116. (Previously presented) The composition of claim 115, wherein said mass tag is a deuterated or non-deuterated amino acid.


117. (Previously presented) The composition of claim 106, wherein said mass tag comprises an element having a characteristic isotope distribution.

118. (Previously presented) The composition of claim 106, wherein said reactive group is an iodoacetyl group.

119. (Previously presented) The composition of claim 106, wherein said photocleavable group comprises amino(ethyl)-2-methoxy-5-nitrophenoxy.

120. (Previously presented) The composition of claim 108, wherein a primary amine group of said polypeptide is modified by treatment with N-succinimidyl S-acetylthioacetate, hydroxylamine, and tris (2-carboxyethyl) phosphine.

121. (Currently amended) A composition comprising having the structure



---

wherein said composition comprises a solid support (1) coupled to a chemical group comprising the cleavable photocleavable functional group, leucine tag and sulfhydryl reactive group shown in Figure 1 said structure, wherein the leucine tag is an isotope tag and wherein said reactive group is covalently linked to a sample molecule.

122. (Previously presented) The composition of claim 121, wherein said sample molecule is selected from the group consisting of a polypeptide, a nucleic acid, a lipid, a second messenger, and a metabolite.

123. (Previously presented) The composition of claim 122, wherein said sample molecule is a polypeptide.

124. (Previously presented) The composition of claim 123, wherein said polypeptide has a modification selected from the group consisting of phosphorylation, glycosylation, ubiquitination, acetylation, palmitoylation, prenylation, sulfation, hydroxylation, and myristylation.

125. (Previously presented) The composition of claim 124, wherein said polypeptide is a phosphopolypeptide.

126. (Previously presented) The composition of claim 121, wherein the solid support is a glass bead.

127. (Previously presented) The composition of claim 123, wherein a primary amine group of said polypeptide is modified by treatment with N-succinimidyl S-acetylthioacetate, hydroxylamine, and tris(2-carboxyethyl)phosphine.

128. (Previously presented) The composition of claim 94, wherein said cleavable group is a photocleavable group, said functional group is a mass tag, and said reactive group reacts with a sulfhydryl group.

129. (Previously presented) The composition of claim 128, wherein the solid support is a glass bead.

130. (Previously presented) The composition of claim 128, wherein said mass tag is an amino acid.

131. (Previously presented) The composition of claim 130, wherein said mass tag is leucine.

132. (Previously presented) The composition of claim 128, wherein said mass tag is selected from the group consisting of a stable isotope tag, an isotope distribution tag, and a charged amino acid.

133. (Previously presented) The composition of claim 132, wherein said mass tag is a stable isotope coded amino acid.


134. (Previously presented) The composition of claim 133, wherein said mass tag is a deuterated or non-deuterated amino acid.

135. (Previously presented) The composition of claim 128, wherein said mass tag comprises an element having a characteristic isotope distribution.

136. (Previously presented) The composition of claim 128, wherein said reactive group is an iodoacetyl group.

137. (Previously presented) The composition of claim 128, wherein said photocleavable group comprises amino(ethyl)-2-methoxy-5-nitrophenoxy.

138. (Currently amended) A composition ~~comprising~~ having the structure



---

wherein said composition comprises a solid support (1) coupled to a chemical group comprising the cleavable photocleavable functional group, leucine tag and sulphydryl reactive group shown in Figure 1 said structure, wherein the leucine tag is an isotope tag.


139. (Previously presented) The composition of claim 138, wherein the solid support is a glass bead.

140. (Previously presented) The composition of claim 138, wherein the leucine tag contains deuterium.

141. (Withdrawn) A method of labeling a sample molecule, comprising contacting a sample with the composition of claim 138 under conditions allowing said sample molecule to covalently bind to said reactive group, and cleaving said cleavable function group, thereby releasing a sample molecule labeled with said tag.



142. (Currently amended) ~~A pair of compounds~~ Two compositions having the structure



each of ~~which comprises~~ said compositions comprising a solid support (1) coupled to a chemical group, said chemical group comprising the ~~cleavable~~ photocleavable functional group, leucine tag and sulfhydryl reactive group shown in Figure 1 said structure, wherein said ~~pair of compounds~~ compositions are differentially isotopically labeled on the leucine tag.

Please add the following new claims.

143. (New) The compositions of claim 142, wherein one of said compositions is differentially isotopically labeled with deuterium on the leucine tag.

144. (New) The compositions of claim 142, wherein one of said compositions is differentially isotopically labeled with <sup>13</sup>C on the leucine tag.